



## **Technical Data Sheet**

| Product Name:      | Zinc Acetate Anhydrous                           |  |
|--------------------|--|--|
| Appearance:        | White Crystal Powder                             |  |
| Solubility:        | Soluble in water & alcohol; insoluble in acetone |  |
| Molecular Formula: | (CH <sub>3</sub> COO) <sub>2</sub> Zn            |  |
| Odor:              | Slight acetic acid odor                          |  |
| Molecular Weight:  | 183.46   |  |
| Specific Gravity:  | 1.74   |  |
| Melting Point:     | 237°C (459°F)                                    |  |
| Boiling Point:     | Decomposes                                       |  |
| CAS No.:           | 557-34-6   |  |
| EINECS No.:        | 209-170-2  |  |
| H.S.Code:          | 2915299090                                       |  |

| Grade                        | Technical Grade          | USP                      |
|------------------------------|--------------------------|--------------------------|
| Appearance                   | White crystalline powder | White crystalline powder |
| Assay (dried basis)          | 98.0 ~ 102.0%            | 98.0 ~ 102.0%            |
| рН                           | 6.0 ~ 7.0                | 6.0 ~ 8.0                |
| Chloride (Cl)                | ≤100ppm                  | ≤50ppm                   |
| Sulfate (SO4)                | ≤100ppm                  | ≤100ppm                  |
| Insoluble matter             | ≤50ppm                   | ≤50ppm                   |
| Iron (Fe)                    | ≤5ppm                    | /                        |
| Lead (Pb)                    | ≤50ppm                   | ≤20ppm                   |
| Arsenic (As)                 | ≤3ppm                    | ≤3ppm                    |
| Alkalies and Alkaline Earths | /                        | ≤0.2%                    |

Packing: 25kgs/bag.

**Storage:** To be stored in cool, dry and ventilated place.

**Uses:** Zinc acetate is used in the pharmaceutical industry, as a buffer in the photographic industry and as a supplement for animal feeds to increase the milk fat production of dairy cattle. It is also used in the production of dye stuffs, as a polymerization catalyst, as a polymer stabilizer, as a flavoring agent, and in the manufacture of hydroxyl oxides, which are used as extractants in hydrometallurgy.